



JOURNALS ON THE WEB

Quick Search in all journals Structure Search

Prous.com

Journals
Home

Methods and Findings
on the Web

Methods and Findings
Information

My
Profile

Contact
Us

Methods and Findings

Methods Find Exp Clin Pharmacol 2000, 22(10): 737
ISSN 0379-0355
Copyright 2000 Prous Science
CCC: 0379-0355
DOI: 10.1358/mf.2000.22.10.802290

Effect of different doses of S-adenosyl-L-methionine on paracetamol hepatotoxicity in a mouse model


Carrasco, R., Perez-Mateo, M., Gutierrez, A., Esteban, A., Mayol, M.J., Cartula, J., Ortiz, P.

This study investigated the hepatoprotective effects of N-acetylcysteine and different doses of S-adenosyl-L-methionine after a single intraperitoneal overdose of paracetamol in mice. Plasma concentrations of paracetamol metabolites were also determined. Female mice (Souris OF1 strain) 16 weeks old and weighing 30 g were fasted for 18 h prior to intraperitoneal (i.p.) administration of 375 mg/kg (2.5 mmol/kg) of paracetamol. Experimental subgroups included mice administered paracetamol only (control group), those given of N-acetylcysteine 1 g/kg (6.13 mmol/kg) i.p. immediately after paracetamol overdose (T0) and 6 h after dosing (T6) and those administered S-adenosyl-L-methionine at doses of 20 mg/kg (0.05 mmol/kg) and 1 g/kg (2.5 mmol/kg) i.p. at T0 and T6. Twenty-four hours after paracetamol overdose, mortality and liver necrosis were significantly lower ($p < 0.01$) in mice treated with 2.5 mmol/kg of S-adenosyl-L-methionine and N-acetylcysteine at T0 as compared with the remaining subgroups. Plasma ALT concentrations were significantly lower ($p < 0.01$) in mice treated with 2.5 mmol/kg of S-adenosyl-L-methionine than in those given N-acetylcysteine. Plasma concentrations of paracetamol metabolites showed an increase in the glucuronide conjugate and a decrease in the mercapturic acid conjugate in N-acetylcysteine-treated mice and an overall decrease in the conjugation pathway without changes in the oxidative pathway in S-adenosyl-L-methionine-treated animals. We conclude that S-adenosyl-L-methionine at doses of 1 g/kg (2.5 mmol/kg) i.p. was equally effective as 1 g/kg (6.13 mmol/kg) N-acetylcysteine for preventing hepatotoxicity after paracetamol overdose in mice. S-adenosyl-L-methionine may be a therapeutic alternative to N-acetylcysteine as an antidote for poisoning with paracetamol.

Full Text: HTML, PDF

Register or sign in

User Name



Alert me to
new issues